arXiv:math/0702254v1 [math.DG] 9 Feb 2007

Singularity Knots of Minimal Surfaces in $\mathbb{R}^4$

Marc Soret & Marina Ville

October 20, 2006

Résumé

We study knots in $\mathbb{S}^3$ obtained by the intersection of a minimal surface in $\mathbb{R}^4$ with a small 3-sphere centered at a branch point. We construct examples of new minimal knots. In particular we show the existence of non-fibered minimal knots. We show that simple minimal knots are either reversible or fully amphicheiral ; this yields an obstruction for a given knot to be an iterated knot of a minimal surface. Properties and invariants of these knots such as the algebraic crossing number of a braid representative and the Alexander polynomial are studied.

1 Introduction

We wish to understand knots associated to specific singularities. Let us first recall what is meant by singularity of a minimal surface. Let D be a disk endowed with a Riemann (complex) structure and let 0 be its center. Let $X : D \mapsto \mathbb{R}^4$ be a conformal harmonic mapping. If $dX(0) = 0$, we say that 0 is a singularity of X. It is then clear that singularities are isolated and correspond to branch points. Let $p = X(0)$. The topology of the singularity at p is entirely determined by a (possibly singular) knot or link which is obtained as follows : intersect M with a small 3-sphere of $\mathbb{R}^4$ around p of radius ϵ. We obtain a curve $K_\epsilon \subset \mathbb{S}^3(\epsilon)$. If p is the image of only one singularity then K_ϵ has only one connected component for ϵ small enough. If $X(D)$ is embedded in a neighborhood of p, then K_ϵ is a smoothly embedded curve and hence is a knot, isotopic to a fixed knot K and $X(D) \cap B(0, \epsilon)$ is topologically a cone over K. For example, a holomorphic complex curve in $\mathbb{R}^4 = \mathbb{C}^2$ is a special case of minimal surface in $\mathbb{R}^4$. When $X(D)$ is locally determined by the equation $F(z,w)=0$, equivalent germs of F at p yield isotopic knots in $\mathbb{S}^3$ (cf. [2]).

1.0.1 Minimal Knots and iterated minimal knots

As is usual in this context, minimal singularities are given by an expansion in terms of z and $\bar{z}$ and one needs to fix some definitions. We remind the reader that a map $X : D \mapsto \mathbb{C}^2$ from the unit disk of $\mathbb{C}$ into $\mathbb{C}^2$ is minimal if and only if X is harmonic with respect to the induced metric on D. However, since a harmonic map from a surface remains harmonic if we change conformally the metric, X is a minimal if X is harmonic with respect to the flat metric on D and if X is conformal ; thus harmonicity means that

$$\Delta_D X = 4\partial_z \partial_{\bar{z}} X = 0. \qquad (\mathcal{H})$$

Let $z = x + iy$; then conformality means that

$$\|\frac{\partial X}{\partial x}\| = \|\frac{\partial X}{\partial y}\|, \quad <\frac{\partial X}{\partial x}, \frac{\partial X}{\partial y}>= 0. \qquad (\mathcal{C})$$

It is straightforward to derive many examples of germs of minimal surfaces :

Definition 1 *Let D be the unit disk in C and let $X : D \longrightarrow \mathbb{R}^4$ and suppose that $X(0) = 0$. The origin 0 is a branch point for X if and only if there is a holomorphic coordinate in D and a coordinate system on $\mathbb{R}^4$ such that X writes in a neighbourhood of 0*

$$z \mapsto \left(Re(z^N) + o(|z|^N),\ Im(z^N) + o(|z|^N),\ o(|z|^N),\ o(|z|^N)\right).$$

We now assume that X is injective. We denote by S_ϵ (resp. B_ϵ) the sphere (resp. ball) in $\mathbf{R}^4$ centered at 0 and of radius ϵ. We put $K_\epsilon = S_\epsilon \cap X(D)$. If ϵ is small enough the following two facts are true
1) $(B_\epsilon, X(D))$ is a cone over (S_ϵ, K_ϵ)
2) all the knots $K_\eta = S_\eta \cap X(D)$ with $\eta \leq \epsilon$ are isotopic. The ensuing knot type is said to be *associated* to the singularity of X at 0.
It follows from the implicit function theorem that there exists a real function r_ϵ such that K_ϵ is parametrized by $X(r_\epsilon(\theta)e^{i\theta})$, with $i\theta$ going through $\mathbb{S}^1$.
If X is a holomorphic map to $\mathbb{C}$, all its singularities are branch points. It is a classical result that the associated knots are iterated torus knots (cf. [2] or

[9]).

Holomorphic curves are a special case of area-minimizing surfaces (Wirtinger inequality) ; these in turn are a special case of minimal surfaces (i.e. conformal harmonic mappings of surfaces).

Micallef and White in [11] have researched branch points of minimal surfaces inside general Riemannian 4-manifolds. They have shown that the area minimizing case is strikingly similar to the holomorphic one : the associated knots are iterated torus knots. In the course of their investigations of general minimal surfaces they showed that other knot types can occur, for example

$$\begin{gathered} D \longrightarrow \mathbb{C}\times\mathbb{C} \\ z \mapsto \left(z^3 + o\left(|z|^3\right), z^4 - \bar{z}^4 + z^5 + \bar{z}^5 + o\left(|z|^5\right)\right) \end{gathered} \tag{1}$$

They noticed that the corresponding minimal knot is the square knot (see fig.3).

Micallef and White left open the following question : can every knot isotopy type can be realized as the knot of a minimal branch point ?

We investigate in the present paper a specific class of knots of branched points of minimal disks. They are given by the following Proposition which is inspired by a remark in [11]

Proposition 1 *Let N, p and q be integers, $p > N$, $q > N$ and let ϕ be a real number. Suppose that the following map*

$$z \mapsto \big(Re(z^N), Im(z^N), Re(e^{i\phi} z^p), Im(z^q)\big)$$

is injective. We denote by $K(N, p, q, \phi)$ the associated knot type.
Then $K(N, p, q, \phi)$ is associated to a branch point of a minimal disk.

Proof. We identify $\mathbb{R}^4$ to $\mathbb{C}^2$ and we look for a map X of the form $X : z \mapsto (z^N + \bar{f}(z), g(z) + \bar{h}(z))$ where f, g and h are holomorphic functions. Such a map is harmonic ; it is moreover conformal if and only if it satisfies equations $(\mathcal{H})$. This translates into

$$f'(z) = -\frac{1}{z^{N-1}} g'(z) h'(z).$$

If $g(z) = z^p + z^q$ and $h(z) = z^p - z^q$, the function f exists and verifies

$$|f(z)| = o(|z|^N).$$

In a neighbourhood of 0, there exists a function $\phi(z)$ such that $\phi(z)^N = 1 + \frac{f(z)}{z^N}$. We put $w = z\phi(z)$; we have $w = z + o(|z|)$. Also $w^N = z^N\phi(z)^N$ and X is of the form

$$X(z) = \left(Re(w^N), Im(w^N), Re[(1+a(w))e^{i\phi}w^p], Im[(1+b(w))w^q]\right)$$

where $a(w)$ and $b(w)$ are $o(1)$. We put

$$X_t(w) = \left(Re(w^N), Im(w^N), Re[(1+ta(w))e^{i\phi}w^p], Im[(1+tb(w))(w^q)]\right).$$

We notice the following obvious

Lemma 1 *Let w and w' be complex numbers, $w \neq w'$, which verify $X_t(w) = X_t(w')$. Then there exists ν a N-th root of 1, $\nu \neq 1$, such that $w = \nu w'$*

In particular $w = w'$. Thus, if we let $w = re^{i\theta}$ we will know that X_t is injective if and only the following maps is injective

$$\tilde{X}_t : D - \{0\} \mapsto \mathbb{R}^4$$

$$\tilde{X}_t : w = re^{i\theta} \mapsto (\cos N\theta, \sin N\theta, Re[(1+ta(w))e^{i\phi}e^{pi\theta}], Im[(1+tb(w))e^{qi\theta}].$$

The value $X_0(re^{i\theta})$ does not depend on r but only on θ, which runs through the compact space $\mathbb{S}^1$. We derive therof the existence of a positive real number $C > 0$ such that for every w, ν, with $w \neq 0$, $\nu^N = 1$ and $\nu \neq 1$,

$$|X_0(w) - X_0(\nu w)| > C.$$

It follows that for $t \in [0,1]$ and w non zero small enough,

$$|X_t(w) - X_t(\nu w)| > \frac{C}{2}.$$

This, together with Lemma 1 above, proves that the X_t's are injective on a small disk around 0. Moreover for ϵ small enough, $X_t(D) \cap S_\epsilon$ constitutes an isotopy between the knots associated to the singularities of X_0 and X. Note that if $q = p$ (and $\phi = 1$ but we will see later that this condition is not necessary), then $K(N,p,q,\phi)$ is a (N,p) torus knot. We call a knot of the type $K(N,p,q,\phi)$ *simple minimal.* The word *simple* comes from the fact that general knots of minimal surfaces can be seen as iterated versions of the $K(N,q,p,\phi)$'s. We plan to devote another paper to these more general knots. For the moment we focus on simple minimal knots.

Notice that simple minimal knots are similar in their expression to Lissajous knots [1], [6], (see also KnotPlot [8] for a generator of Lissajous knots), and is a good help in the understanding of the latter ones. Still their properties are quite different as we shall see.

Let us give a brief outline of the paper.

In section 1, we give a general description of the properties of simple minimal knots which are parametrized by three numbers and a phase $K(N,p,q,\phi)$; we then give a geometric interpretation of N,p,q and description of the braid representation that is naturally attached to them.

In section 2, the minimal braid representation is studied in more detail and we show that simple knots are invariant by a change of phase. Minimal knots will be denoted by $K(N,p,q)$.

In section 3, we study the symmetries of minimal knots. Recall that there are two natural symmetries among knots that are involutions ; the mirror symmetry s_m, (symmetry of a knot with respect to a orientation reversing symmetry of $\mathbb{S}^3$) and the inversion of a knot s_i which maps a knot to the same knot but with the reverse orientation. K is invertible if it is invariant by s_i and amphicheiral if it has some invariance with respect to s_m Results of section 2 and easy computations yields

Theorem 1 *A simple minimal knot is either reversible or fully amphicheiral. More precisely :*

1. *All knots $K(N,p,q)$ are strongly invertible.*
2. *If $p+q$ is odd, then (N,p,q) is strongly + amphicheiral :*
3. *If N is even and $p+q$ is even or if N is odd and p and q even, then $K(N,p,q)$ is periodic of order two. The S^1 curve by the involution invariant has a linking number with $K(N,p,q)$ equal to N.*

By a result of [5] and [10], and similarly to the Lissajous knots (cf. [1]), these symmetries yields properties on the Arf invariant and Alexander polynomial that are described in the section.

Section 3 leads to section 4 where we show the existence of knots that can not be realized as simple or iterated knots of minimal singularities. These are the negative amphichireal or chiral knots.

Theorem 2 *A negative amphicheiral or chiral knot can not be the knot of a simple minimal or iterated minimal knot*

The first candidate in the Rolfsen classification is the knot 8_{17} which is the first negative amphicheiral knot, and 9_{32} which is the first chiral knot.

We do not yet know if these chiral knots can be realized as cable knots of minimal singular knots. This question will be treated elsewhere.

Section 6 is devoted to the algebraic crossing number of the natural braid representation (see Section 1) of simple minimal knots. We will give an upperbound of this number.

With the help of KnotTheory and KnotPlot, we describe in section 7 some examples of minimal knots with their minimal braid representation and decomposition into prime knots. This allows us to investigate the fibration of minimal knots. Let us first recall a few definitions.

Definition 1 *K is fibered if $S^3 \backslash K$ is fibered over S^1 : there is a differentiable mapping $\phi : \mathbb{S}^3 \setminus K \mapsto S^1$ which defines a fiber bundle ; the fiber $\phi^{-1}(e^{it})$ is the interior of a compact orientable differentiable surface with boundary K.*

(Notice that fibered knots have an algebraic characterization : the commutator of the knot group $\pi_1(\mathbb{S}^3 \setminus K)$ is finitely generated . [4]) Knots of holomorphic curves singularities are always fibered ; if the surface is given locally by the equation $F(z, w) = 0$, then the (Milnor) fibration is simply given by

$$\phi(z, w) = \frac{F(z, w)}{|F(z, w)|}$$

restricted to a sphere of sufficiently small radius $\mathbb{S}^3(p, \epsilon)$.

Thus all torus knots are fibered.

The fibration yields a monodromy mapping h and a gluing map θ where $\mathbb{S}^3 \setminus K \equiv \frac{I \times \mathcal{F}}{(x,0) \sim (\theta(x),1)}$, where $\mathcal{F}$ is the fiber of the fibration, ie the Seifert surface spanning K.

The monodromy map gives a way to compute the Alexander polynomial, another knot invariant :

$$P(x) = det(h^* - xId)$$

where $h^* : H^1(M, \mathbb{R}) \mapsto H^1(M, \mathbb{R})$.

In particular if the highest order term of P is different from ±1 then the knot is can not be fibered.

We construct some examples of minimal braids with three, four or five strands and identify them. In particular :

Theorem 3 *The knot* 9_{46} *(see fig. 13) is a prime knot which is not fibered and still is a minimal singularity knot ; the minimal surface is locally given by*

$$z \mapsto \left(z^4 + o(|z|^4),\ z^{13} + \bar{z}^{13} + z^5 - \bar{z}^5 + o(|z|^{13})\right).$$

We conjecture that the simple knots $K(N,p,q)$ with N odd are fibered.

We are grateful to Harold Rosenberg for introducing us to the problem of fibration of minimal singularities. We thank Joan Birman for her suggestions and Dror bar Natan for his help and diligency.

2 Definition of a minimal knot and description of its minimal braid

2.0.2 Germ of a minimal singularity

At a branch point p, a minimal surface M still has a tangent plane $\mathbf{TM_p}$ which is a real plane in $\mathbb{C}^2$. The ambiant space is thus split into $\mathbf{TM_p}$ and the normal plane $\mathbf{NM_p}$. M is a multi-valued minimal graph over a subdomain of $\mathbf{TM_p}$ locally given by (proposition 1)

$$z \mapsto \left(z^N,\ a(z^p + \bar{z}^p) + b(z^q - \bar{z}^q)\right)$$

$a, b \in \mathbb{C}$. Let us intersect M with a 3-cylinder $S_\epsilon \times \mathbf{NM_p} \subset \mathbf{TM_p} \times \mathbf{NM_p}$, perpendicular to $\mathbf{TM_p}$; S_ϵ is a small circle of radius ϵ in $\mathbf{TM_p}$.The intersection a knot parametrized by θ running along $\mathbb{S}^1(\epsilon)$ N times. As ϵ tends to zero, this knot converges clearly to the equator of the corresponding 3-sphere $\mathbb{S}^3(0,\epsilon)$ hence K can be equally viewed as a knot of the 3-sphere.

This knot can be expressed in terms of circular functions as follows :

$$\left\{ \begin{array}{c} K : \mathbb{S}^1(0,\epsilon) \longrightarrow \mathbb{R}^4 \\ \theta \mapsto (\cos N\theta,\ \sin N\theta,\ \cos{(p\theta + \phi_p)},\ \sin{(q\theta + \phi_q)}) \end{array} \right. \tag{2}$$

Changing θ into $\theta + \alpha$, we may arrange that one of the two phases is zero. We choose the following parametrization

$$K(N, p, , q, \phi) : \theta \mapsto (\cos N\theta, \sin N\theta, \cos(p\theta + \phi), \sin(q\theta))$$

We parametrize $\mathbb{S}^1$ by $t \in [0,1]$ where $\theta = 2\pi t$.

Definition 2 *A simple minimal knot, is isotopic to the curve given by the one-to-one parametrization*

$$B : [0,1] \longrightarrow \mathbf{Cyl} \subset \mathbb{R}^4$$

$$t \mapsto (\cos 2\pi Nt, \sin 2\pi Nt, \cos(2\pi pt + \phi), \sin 2\pi qt)$$

We denote $B([0,1])$ *by* $K(N,p,q,\phi)$

These knots are similar to another type of knots, known as Lissajous knots because the coordinates of the knot are circular functions of possibly unequal frequencies.

2.0.3 Lissajous Knots

It is interesting to note another connection between Minimal knots and Lissajous knots. Let us first recall how it was defined in ([1] or [6])

Definition 3 *A Lissajous knot is a curve parametrized one-to-one by*

$$L : [0,1] \longrightarrow \mathbb{R}^3$$
$$t \mapsto (\cos 2\pi Nt, \cos(2\pi pt + \phi_1), \cos(2\pi qt + \phi_2))$$

Notice that these knots are parametrized by 5 variables instead of 4 for simple minimal knots.

Lemma 2 *The projection into a vertical 3-plane of a minimal knot is a Lissajoux knot. If the minimal knot is of type* $K(N,p,q)$ *then the corresponding Lissajous knot has a braid representation of 2N strands induced by the braid representation of* $K(N,p,q)$*.*

This is a direct consequence of the definition of a Lissajoux knot.

2.0.4 The minimal braid representation of a simple minimal knot

The graph of the function $B : S_\epsilon \mapsto \mathbb{C}$ defined by $K(N,p,q,\phi)$, is a braid of N strands : there are N functions defined on the open circle or $[0,1[$ i.e. $k = 0, ..., N-1$ such that

$$B_k : [0,1] \longrightarrow \mathbb{C}$$



FIG. 1 – braid and braid diagram of K(3,4,4)

$$B_k(t) = \cos\left(\frac{2\pi p}{N}(t+k) + \phi_p\right) + i \sin\left(\frac{2\pi q}{N}(t+k)\right)$$

Reciprocally we reconstruct the knot $K(N,p,q,\phi)$ by closing the braid : we connect the kth strand to the (k+1)th strand as follows : $B_k(1) = B_{k+1}(0)$ for $k = 0, ..., N-1$.

We denote this braid by $B(N,p,q,\phi)$. We then project the braid onto any $\mathbb{R} \subset \mathbb{C}$.

Definition 4 *A braid diagram is the graph of the functions $\pi_D \circ B$ where π_D is the projection of $\mathbb{C}$ onto a line D of $\mathbb{C}$.*

If we choose to project the braid onto the y-component of $\mathbb{C}$, $(z = x+iy)$, we obtain a braid diagram that depends only on N and q. We denote it by $K^{\perp}(N,q)$. This braid diagram consists of N graphs of functions

$$h_k = ImB_k : [0,1] \longrightarrow \mathbb{R},$$

$k = 0, \cdots, N-1$. Note that all the $K^{\perp}(N,q)$ are identical to the braid diagram of the (N,q)-torus knot. To regain the braid from the braid diagram, we need to know which strand is above or below at each intersection of any two strands h_k and h_l. We will first determine the value of the parameter t for each crossing and will determine which strand is above at each crossing.

2.1 Crossing locus of the braid diagram $K^{\perp}(N,q)$

Let us compute the parameter values t at each intersection points of the braid diagram $K^{\perp}(N,q)$.

Choose $k < l$, $0 \leq k \leq N-1$, $t \in [0,1[$; we study the intersection of the strands h_k et h_l ; the crossings correspond to the different values t for which

$$h_k(t) - h_l(t) = 0.$$

It turns out that for an even number N of strands the parameter value $t=0$ corresponds to a value for which two strands h_{k_1} and h_{k_2} intersect ; we thus introduce a small positive ϵ such that ϵ does not parametrize an intersection of two strands ; we study instead the braid diagram on the new interval $t \in [\epsilon, 1+\epsilon[$.

Lemma 1 *Let $K(N,p,q,\phi)$ be a simple minimal knot. Its braid diagram $K^{\perp}(N,q)$ consists of N strands ; its crossing number is $q(N-1)$. Furthermore the strings h_k and h_l, $k<l$, $k=0,...,N-1$, meet at points $(t_{n,k,l}, h(t_{n,k,l}))$, $t \in [\epsilon, 1+\epsilon[$ where*

$$t(n,k,l) := \frac{N(2n+1)}{4q} - \frac{k+l}{2};$$

n is any integer such that

$$\frac{2q\epsilon}{N} + \frac{q(k+l)}{N} - \frac{1}{2} \leq n < \frac{2q\epsilon}{N} + \frac{q(k+l)}{N} - \frac{1}{2} + \frac{2q}{N}.$$

Proof.

$$h_k(t) - h_l(t) = 0$$

$$\Im B_k(\theta) - \Im B_l(\theta) = \sin 2\pi \frac{q}{N}(t+k) - \sin 2\pi \frac{q}{N}(t+l)$$

using

$$\sin a - \sin b = 2\cos(a+b/2)\sin(a-b/2)$$

we obtain

$$\Im B_k(\theta) - \Im B_l(\theta) = 2\cos\left(2\pi \frac{qt}{N} + \pi \frac{q(k+l)}{N}\right) \sin \pi \frac{q(k-l)}{N} \qquad (3)$$

It follows from (3), that t parametrizes a double point if the first cosine factor is zero that is

$$2\pi \frac{qt}{N} + \pi \frac{q(k+l)}{N} = \pi \frac{2n+1}{2}$$

or

$$t = \frac{N(2n+1)}{4q} - \frac{k+l}{2}.$$

Hence the number of intersections of $K^{\perp}(N,q)$ is given by the number of integers n such that

$$\epsilon \leq \frac{N(2n+1)}{4q} - \frac{k+l}{2} < 1 + \epsilon$$

$$\epsilon + \frac{k+l}{2} \leq \frac{N(2n+1)}{4q} < 1 + \frac{k+l}{2} + \epsilon$$

$$\frac{4q}{N}\epsilon + \frac{2q(k+l)}{N} \leq (2n+1) < \frac{4q}{N} + \frac{2q(k+l)}{N} + \frac{4q}{N}\epsilon$$

$$\frac{2q}{N}\epsilon + \frac{q(k+l)}{N} - \frac{1}{2} \leq n < \frac{2q}{N} + \frac{q(k+l)}{N} - \frac{1}{2} + \frac{2q}{N}\epsilon$$

□

Example 1 : N=2 The integer n that parametrizes the solutions verifies

$$q \leq (2n+1) < 3q$$

$$\frac{q-1}{2} \leq n < \frac{q-1}{2} + q$$

It follows that the braid diagram has q crossing points.

2.1.1 Regularity of $K(N,p,q,\phi)$

We return to the minimal knot and its naturally associated braid. The braid may be be singular if some strands intersect : if this is the case, $B_k(t) = B_l(t)$ for some k, l. and t. To solve this equation, we need the following lemma complementary to lemma 1

Lemma 2 *the projections $\Re B_k$ and $\Re B_l$ $k < l$ of $K(N,p,q,2\pi\varphi)$ meet at points*

$$t_{n,k,l} := \frac{mN}{2p} - \frac{N\varphi}{2p} - \frac{k+l}{2}$$

where m is any integer such that

$$\frac{p}{N}(k+l) + 2\varphi \leq n < \frac{p}{N}(k+l) + 2\varphi + 2\frac{p}{N}$$

Proof. Let $\phi = 2\pi\varphi$

$$\cos\left(2\pi\frac{p}{N}(t+k)+\phi\right)-\cos\left(2\pi\frac{p}{N}(t+l)+\phi\right)=0$$

if and only if

$$\sin\left(2\pi\frac{p}{N}\left(t+\frac{k+l}{2}\right)+\phi\right).\sin\left(2\pi\frac{p}{N}(k-l)\right)=0$$

$$2\pi\frac{p}{N}\left(t+\frac{k+l}{2}\right)+\phi=\pi m$$

$$t=\frac{mN}{2p}-\frac{N\varphi}{2p}-\frac{k+l}{2}$$

$$t\in[\epsilon,1+\epsilon]$$

iff

$$\epsilon\frac{2p}{N}+\frac{p}{N}(k+l)+2\varphi\leq m<\frac{p}{N}(k+l)+2\varphi+2\frac{p}{N}+\epsilon\frac{2p}{N}$$

Hence the simple knot has selfintersection, if there is a $t \in [\epsilon, 1+\epsilon[$ that satisfies the hypothesis of lemma 1 and lemma 2 ; we conclude that the braid is singular if there are integers m and m' such that

$$(2m+1)p-2qm'=-4\varphi q$$

On the other hand if N and p or N and q are not coprime.then $K(N,p,q,\phi)$ is always singular : indeed if $N = aN',\ p = ap'$, then the substitution $t \mapsto t+\frac{k}{a}$ doesn't alter the first 3 coordinates of the knot : it suffices to verify that that we may choose $\epsilon < t < 1+\epsilon$ such that the last coordinate has the same value for t and $t+\frac{k}{a}$.

Proposition 1 *Let $K(N,p,q,\phi)$ a minimal knot ; if $N\wedge p = N\wedge q = 1$, then for almost all ϕ $K(N,p,q,\phi)$ is regular. $K(N,p,q,\phi)$ is singular for a finite number of ϕ ; and these ϕ's are all of the form $\phi = 2\pi\alpha$, where α is rational.*

Example. The knots $K(N,p,p,\phi)$ are the torus knots (N,p). The knot projects on the first component of $\mathbb{C}^2$ onto a circle and on the second component of $\mathbb{C}^2$ onto an ellipse if $\phi \neq 0$ or a segment if $\phi = 0$. The "knot" is clearly singular for its value. We will show in section 3 that, surprisingly, a minimal knot does not change its type as the phase varies. This is a striking difference with the Lissajous knots where a suitable conjugate variation of the two phases may change the knot ([1]).



-1



+1

2.2 The y-coordinate of a crossing point

Let $t(m,k,l)$ be a crossing point between the k-th and l-th strands as given by Lemma 1. The y-coordinate of the corresponding point of the braid, i.e. the height of the braid diagram is given by

$$h_k(t) = h_l(t) = y(t,k,l) = \sin\frac{2\pi}{N}q(t{+}k) = (-1)^m \sin\left(\frac{\pi}{N}q\,(k-l) + \frac{\pi}{2}\right) \quad (4)$$

2.3 Sign of the crossing points

If t is a crossing point between the k-th and l-th strand, its sign $S(t,k,l)$ is given by combining the following two pieces of data :

1. which strand is in front of the other one
2. which strand goes upwards.

(1) is given by the sign of the difference $ReB_k(t) - ReB_l(t)$, that is the sign of the difference in the x-coordinates

$$\cos\frac{2\pi}{N}q(t+k+\phi) - \cos\frac{2\pi}{N}q(t+l+\phi)$$

For (2), we notice that the k-th strand is going upwards if the derivative of the function $x \mapsto \sin\frac{2\pi}{N}qx$ at $t+k$ is positive (in which case its derivative at $t+l$ is negative). It follows that the sign is given by the product

$$S(t,k,l) = -[\cos q(t+k) - \cos q(t+l)][\cos p(t+k+\phi) - \cos p(t+l+\phi)].$$

We remind the reader that a difference of cosines can be written as a product of sines and derive

$$S(t,k,l) = -4\sin\frac{2\pi}{N}q(t{+}\frac{k+l}{2})\sin\frac{2\pi}{N}p(t{+}\frac{k+l}{2}{+}\phi)\sin\frac{\pi}{N}p(k{-}l)\sin\frac{\pi}{N}q(k-l).$$

We now suppose that t is of the form $t = t(m,k,l)$ as given by lemma 2. We derive

$$\sin\frac{2\pi}{N}q(t+\frac{k+l}{2}) = \sin(\frac{\pi}{2} + m\pi) = (-1)^m$$

$$\sin\frac{2\pi}{N}p(t+\frac{k+l}{2}+\phi) = \sin\frac{2\pi}{N}p\left(\phi + \frac{N}{4q}\,(1+2m)\right).$$

Notations. If x is a real number we denote its integral part by $[x]$; if n is an integer, we denote by $P(n) \in \mathbb{Z}_2$ its congruence modulo 2.
With these notations, and with the notations of lemma (1) last computations give the following lemma

Lemma 3 Crossing sign formula
The sign at a crossing point of the braid diagram $K(N,q)^{\perp}$ associated to the simple knot $K(N,p,q,\phi)$ and parametrized by $t_{m,k,l} \in [\epsilon, 1+\epsilon[$, where $k<l$, $k=0,...,N-1$, and $m \in \mathbb{N} \cap [\frac{2q\epsilon}{N} + \frac{q(k+l)}{N} - \frac{1}{2}, \frac{2q\epsilon}{N} + \frac{q(k+l)}{N} - \frac{1}{2} + \frac{2q}{N}[$, is given by the formula

$$s(k,l,m,\phi) = 1 + T(k,l) + P(m) + R(m,\phi) \tag{5}$$

with respectively

$$T(k,l) = P\left([q\left(\frac{k-l}{N}\right)]\right) + P\left([p\left(\frac{k-l}{N}\right)]\right)$$

(the sum is taken in $\mathbb{Z}_2$) and

$$R(m,\phi) = P\left([\frac{2}{N}p(\phi + \frac{N}{4q}(1+2m))]\right).$$

2.4 An expression of a minimal braid

We recall here the definition of the braid groups in terms of generators and relations.
The braid group $\mathbf{B}_N$ is generated by $N-1$ elements, $\sigma_1, ..., \sigma_{N-1}$ subject to the following relations

$$\forall i, \quad \sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1}$$

$$\forall i,j \text{ with } |i-j| \geq 2, \quad \sigma_i \sigma_j = \sigma_j \sigma_i.$$

Since the braid $B(N,p,q,\phi)$ has $(N-1)q$ crossing points, we can write it as a product of $(N-1)q$ σ_i's or σ_i^{-1}'s. It is straightforward to derive such an expression; however because of the above-mentioned relations there are several such expressions and we need to specify exactly which one we take. In other words we will specify one representative of a preimage of $B(N,p,q,\phi)$ in the free group $\mathbf{F}_{N-1}$ on the $N-1$ generators σ_i's.

Up to the sign of the crossing points, the braid is the same as for a $(N-1)q$-torus knot, thus we aim for a representation which differs from the standard representation of the torus knot only by the signs of the terms. Each crossing point $C(t,k,l)$ is the data of coordinate t and a pair of strands k and l. We number then as

$$C_1(t_1,\{k,l\}_1),...,C_{q(N-1)}(t_{q(N-1)},\{k,l\}_{q(N-1)}).$$

We require the following ordering conditions

1. if $u<v$ then $t_u \leq t_v$
2. the first $\frac{(N-1)N}{2}$ crossing points are ordered by lexicographical order on $t, k+l, k-l$
3. if $u+\frac{N(N-1)}{2} \leq q(N-1)$, then $\{k,l\}_{u+\frac{N(N-1)}{2}} = \{k,l\}_u$.

Each crossing point $C_i(t_i,\{k,l\}_i)$ has a sign $\epsilon(i) \in \{-1,1\}$ and a y-coordinate $y(C_i)$, (cf. 4). We point out that the value in (cf. 4) takes $N-1$ different values $y_1 > y_2 > ... > y_{N-1}$.
We define $n(i)$ by

$$n(i) = s \text{ if and only if } y(C_i) = y_s.$$

We derive a word in $\mathbf{F}_{N-1}$

$$b_{N,q,p,\phi} = \prod_{i=1}^{(N-1)q} \sigma_{n(i)}^{\epsilon(i)}.$$

Its image in $\mathbf{B}_N$ is a representative of $B(N,p,q,\phi)$.
Since we will be dealing with words of length $q(N-1)$ we introduce, for an integer m, the sets

$$\mathbb{N}_m = \{n \in \mathbb{N} \mid 1 \leq n \leq m\}.$$

Thus the word $b_{N,q,p,\phi}$ is given by the data of the maps

$$n : \mathbb{N}_{q(N-1)} \longrightarrow \mathbb{R}^{N-1}, \quad \epsilon : \mathbb{N}_{q(N-1)} \longrightarrow \mathbb{Z}_2.$$

3 (Non) dependence of the knot type on the phase ϕ

Proposition 2 *Let N,p,q be integers as above and let ϕ and ϕ' be two elements of $[0,2\pi]$. Then the knots $K(N,p,q,\phi)$ and $K(N,p,q,\phi')$ - or $K(N,q,p,\phi)$ and the mirror image of $K(N,q,p,\phi')$- can be represented by conjugate braids.*

Corollary 1 *Up to taking a mirror image, the isotopy type of the knot* $K(N, q, p, \phi)$ *does not depend on the phase* ϕ.

Lemma 3 *It is enough to prove the Proposition in the case*

$$\phi' = \phi + \frac{N}{2}\left(\frac{A}{p} + \frac{B}{q}\right)$$

for two integers A *and* B.

To prove the lemma we introduce

Definition 5 *An element* $\phi \in [0, 2\pi]$ *is said to be a* ***critical phase*** *if there exists a crossing point* $t = t(m, k, l)$ *so that*

$$\cos\frac{2\pi}{N}p(t + k + \phi) = \cos\frac{2\pi}{N}p(t + l + \phi).$$

In that case $K(N, p, q, \phi)$ has self-intersection at $t(m, k, l)$. This translates into the existence of an integer s such that

$$\frac{2\pi}{N}(t + k + \phi)p = -\frac{2\pi}{N}(t + l + \phi)p + 2\pi s$$

hence

$$\phi = \frac{sN}{2p} - \frac{N}{4q}(1 + 2m) \tag{6}$$

The braid $B_{N,q,p,\phi}$ is defined if and only if the number ϕ is not a critical phase.

The following is obvious

Lemma 4 *Let* ϕ_1, ϕ_2 *be two numbers,* $\phi_1 < \phi_2$ *and suppose that there is no critical phase in the interval* $[\phi_1, \phi_2]$. *Then the braids* B_{N,q,p,ϕ_1} *and* B_{N,q,p,ϕ_2} *are identical.*

We have

Lemma 5 *Let* ϕ_1 *and* ϕ_2 *be two critical phases. Then there exist two integers* A *and* B *such that*

$$\phi_1 - \phi_2 = \frac{N}{2}\left(\frac{A}{p} + \frac{B}{q}\right).$$

Proof We write ϕ_1 and ϕ_2 as in (6) in terms of s_1, m_1 and s_2, m_2. We have

$$\phi_1 - \phi_2 = (s_1 - s_2)\frac{N}{2p} - \frac{N}{2q}(m_1 - m_2)$$

and the lemma follows.
We order the critical phases between 0 and 2π and denote them respectively

$$\phi_0,\ \phi_1,\ \cdots, \phi_M.$$

We assume that $\phi_u < \phi < \phi_{u+1}$ and $\phi_v < \phi' < \phi_{v+1}$ for some u, v, with $0 \leq u, v \leq M$. It follows from the lemma that $\phi_u - \phi_v = X$ with $X = \frac{N}{2}(\frac{A}{p} + \frac{B}{q})$ for some integers A, B. Thus $\phi + X$ belongs to the interval $[\phi_v, \phi_{v+1}]$ (like ϕ' does). Hence the braids $B_{N,q,p,\phi'}$ and $B_{N,q,p,\phi+\frac{N}{2}(\frac{A}{p}+\frac{B}{q})}$ are the same. $\square$

Going back to the formula for the signs of the crossing points, a straightforward computation yields

Lemma 6 *Let N, q, p, ϕ be as above and let A and B be two integers. Then*

$$R\left(m, \phi + \frac{N}{2}(\frac{A}{p} + \frac{B}{q})\right) = P(A) + R(m + B, \phi).$$

In view of 3, Lemma 5 solves the problem if $B = 0$: $B_{N,q,p,\phi}$ is the same (resp. mirror image) of $B_{N,p,q,\phi+\frac{AN}{2p}}$ if A is even (resp. odd). Hence Lemma 6 will be proven (by induction) once we have shown that $K_{N,q,p,\phi}$ and the mirror image of $K_{N,q,p,\phi+\frac{N}{2q}}$ can be represented by conjugate braids. More precisely we will show

Lemma 7 *Let $\Phi : B_N \longrightarrow B_N$ be the involutive isomorphism defined by*

$$\forall i, \quad 1 \leq ... \leq N-1, \quad \Phi(\sigma_i) = \sigma_{N-i}.$$

1. *$\Phi(B_{N,q,p,\phi})$ is a braid which represents $K_{N,q,p,\phi}$.*
2. *The mirror image of $\Phi(B_{N,q,p,\phi})$ and $B_{N,q,p,\phi+\frac{N}{2q}}$ are conjugate braids.*

Proof 1. is obvious and we set out to prove 2.
We consider the canonical representative $b(N, p, q, \phi + \frac{N}{2q})$ which we have described in the previous paragraph. Then

$$b(N, p, q, \phi + \frac{N}{2q}) = \prod_{i=1}^{q(N-1)} \sigma_n(i)^{\eta(i)}.$$

We then write a word in $\mathbf{F}_{N-1}$ which represents the mirror image of $\Phi(B_{N,q,p,\phi})$: we take $b_{N,q,p,\phi}$, change every exponent into its inverse and replace every σ_i by the corresponding σ_{N-i} ; we derive the word

$$\tilde{b}_{N,q,p,\phi} = \prod_{i=1}^{(N-1)q} \sigma_{N-n(i)}^{-\epsilon(i)}.$$

Both $b_{N,q,p,\phi+\frac{N}{2q}}$ and $\tilde{b}_{N,q,p,\phi}$ are words of length $q(N-1)$ in the free group F_{N-1} ; part 2) of the lemma will derive from

Sublemma 1 *The words $b_{N,q,p,\phi+\frac{N}{2q}}$ and $\tilde{b}_{N,q,p,\phi}$ are a circular permutation of one another.*

The permutation in question is given by

$$\gamma : \mathbb{N}_{q(N-1)} \longrightarrow \mathbb{N}_{q(N-1)}$$

$$(1,2,...,q(N-1)) \mapsto \left(1+\frac{N(N-1)}{2}, 2+\frac{N(N-1)}{2}, \cdots, \frac{N(N-1)}{2}-1, \frac{N(N-1)}{2}\right).$$

In view of the notations we have chosen, proving the Sublemma means proving the following two facts for every $i \in \mathbb{N}_{q(N-1)}$
(S1) $n(i) = N - n(\gamma(i))$,
(S2) $\eta(i) = 1 + \epsilon(\gamma(i))$.

1. **First case** $i + \frac{N(N-1)}{2} \leq (N-1)q$.

   Conditions 1) and 3)of section 2.4 on the ordering of the crossing points $C_s(t_s, \{k,l\}_s)$ ensure that C_i and $C_{i+\frac{N(N-1)}{2}}$ are consecutive (in $[0,1]$) crossing points of the pair of strands $\{k,l\}_i$, hence

   $$\{k,l\}_{\gamma(i)} = \{k,l\}_i.$$

   We also derive that, if $t_i = t(m_i,k_i,l_i) = -\frac{k_i+l_i}{2} + \frac{N}{4q}(1+2m_i)$, then

   $$t_{i+\frac{N(N-1)}{2}} = -\frac{k_i+l_i}{2} + \frac{N}{4q}\left(1+2(m_i+1)\right).$$

   Thus $t(m_i,k_i,l_i) - t(m_i+1,k_i,l_i) = \frac{N}{2q}$. Hence

$$y(C_i) + y(C_{i+1}) = \sin\frac{2\pi}{N}q(t_i) + \sin\frac{2\pi}{N}q(t_i + \frac{N}{2q}) = 0$$

It follows that

$$n(\gamma(i)) = N - n(i)$$

This proves (S1).
The number $\eta(i)$ gives us the sign of the crossing point $C_i(t_i, k_i, l_i)$ of the braid $B(N, p, q, \phi + \frac{N}{2q})$. It follows from 3 and Lemma 4 that

$$\eta(i) = 1 + T(k_i, l_i) + P(m_i) + R(m_i + 1, \phi) = T(k_i, l_i) + P(m_i + 1) + R(m_i + 1, \phi).$$

We derive from the considerations above that the right-hand side is equal to $1 + \epsilon(\gamma(i))$. This proves (S2).

2. **Second case**. $i + \frac{N(N-1)}{2} > q(N-1)$
Then C_i is the last crossing point of the strands $\{k, l\}_i$; thus the reasonning of the previous case does not work word for word ; however we will establish a $1-1$ correspondence between the last $\frac{N(N-1)}{2}$ crossing points of the braid and the first $\frac{N(N-1)}{2}$ ones. We put

$$t(m_i + 1, k_i, l_i) = -\frac{k_i + l_i}{2} + \frac{N}{4q}\left(1 + 2(m_i + 1)\right) > 1$$

Please note the slight abuse of notation : as we said, $t(m_i + 1, k_i, l_i)$ is NOT a crossing point for the k_i-th and l_i-th strands. We have

$$0 < t(m_i + 1, k_i, l_i) - 1 = -\frac{k_i + l_i + 2}{2} + \frac{N}{4q}\left(1 + 2(m_i + 1)\right) < 1.$$

We recognize the formula given for a crossing point of the braid.
It is clear that the $t(m_i + 1, k_i, l_i) - 1$ occur in the same order (in the interval $[0, 1]$) as the $t(m_i, k_i, l_i)$'s.
We need to distinguish two subcases.

   (a) **First subcase of second case**. $k_i < l_i < N - 1$.
   Then $t(m_i + 1, k_i, l_i) - 1$ is a crossing point for the strands $k_i + 1, l_i + 1$. It is the first crossing point for that pair of strands and writes in our notation

$$t(m_i + 1, k_i + 1, l_i + 1).$$

We have $k_\gamma(i) = k_i{+}1,\ l_\gamma(i) = l_i{+}1$ and in terms of y coordinates,

$$y(C(m_i{+}1, k_i{+}1, l_i{+}1)) = (-1)^{m_i+1}\sin[\frac{\pi}{N}q(k_i{-}l){+}\frac{\pi}{2}] = -y(C(t_i, k_i, l_i)).$$

Thus $(S1)$ is true.
Since $T(k_i+1, l_i+1) = T(k_i, l_i)$, the sign of $t(m_i+1, k_i+1, l_i+1)$ as a crossing point of the k_i+1-th and l_i+1-th strands of $B_{N,q,p,\phi}$ is given by

$$\epsilon(\gamma(i)) = 1 + T(k_i, l_i) + P(m_i + 1) + R(m_i + 1, \phi)$$

$$= 1 + T(k_i, l_i) + 1 + P(m_i) + R(m_i, \phi + \frac{N}{2q}) = 1 + \eta(i).$$

(b) **Second subcase of second case**. $k_i < l_i = N - 1$.
That is, we are considering a point of the form $t(m_i, N-1, l_i)$. We have

$$t(m_i + 1, N - 1, l_i) - 1 = -\frac{N + l + 1}{2} + \frac{N}{4q}\left(1 + 2(m_i + 1)\right)$$

$$= -\frac{l + 1}{2} + \frac{N}{4q}\left(1 + 2(m_i + 1 - q)\right).$$

Thus $t(m_i{+}1, N{-}1, l_i){-}1$ is a crossing point for the l_i-th and 0-th strands. According to our notations it writes $t(m_i+1-q, 0, l_i+1)$. We have

$$\{k, l\}_{\gamma(i)} = \{N - 1, l_{i+1}\}$$

$$y(C(m + 1 - q, 0, l + 1)) = (-1)^{m+1-q}\sin\left(\frac{\pi}{N}q(l + 1) + \frac{\pi}{2}\right)$$

On the other hand,

$$y(C(m_i, N - 1, l_i)) = (-1)^m \sin[\frac{\pi}{N}q(N - 1 - l) + \frac{\pi}{2}]$$

$$= (-1)^m \sin[(q + 1)\pi - \frac{\pi}{N}(l + 1) - \frac{\pi}{2}] = y(C(m_i + 1 - q, 0, l_i + 1))$$

This proves (S1) ; to investigate the sign of $t(m_i + 1 - q, 0, l_i + 1)$ we need the following identities

$$T(N - 1 - l_i, 0) = P(p) + P(q) + T(l_i + 1, 0)$$

$$\begin{aligned} P(m_i+1-q) &= P(q)+P(m_i)+1 \\ R(m_i+1-q,\phi) &= P([\frac{2\pi}{N}(\phi+\frac{N}{4q}(1+2m_i)+\frac{N}{2q}-\frac{N}{2}]) \\ &= R(m_i,\phi+\frac{N}{2q})+P(p). \end{aligned}$$

Summing all these terms, we see that

$$s(m_i+1-q,l+1,0,\phi)=1+s(m_i,N-1,l,\phi+\frac{N}{2q}).$$

which proves $(S2)$ in this case.

This concludes the proof of the lemma and hence, of the Proposition.

4 Symmetries of the knot

Let us recall some terminology on knots symmetries : There are two natural symmetries among knots that are involutions ; the mirror symmetry s_m, (symmetry of a knot with respect to a orientation reversing diffeomorphism of $\mathbb{S}^3$) and the inversion of a knot s_i which yields the same knot but with the reverse orientation. K is *invertible* if it is invariant by s_i and *amphicheiral* if it has some invariance with respect to s_m ; more precisely, each type of invariance is given a name and we recall them here for clarity's sake.

1. If $s_i(K)=K$ (equal means isotopic) only, then K is reversible ;
2. if $s_i(K)=K$ and $s_m(K)=K$ then K is fully amphicheiral ;
3. If $s_m(K)=K$ only, then K is (positive) amphicheiral ;
4. if $s_m\circ s_i(K)=K$ only, then K is negative amphicheiral.
5. If K has none of these symmetries then K is chiral.

The knot is said to be strongly symmetric if it is symmetric with respect to an ambiant isometry of $\mathbb{S}^3$.

Results of section 2 and easy computations yields

Theorem 4 *A simple minimal knot is either reversible or fully amphicheiral. More precisely :*

1. *All knots $K(N,p,q)$ are strongly invertible.*

2. *Furthermore, if $p+q$ is odd, then (N,p,q) is strongly fully amphicheiral :*
3. *if N is even and $p+q$ is even or if N is odd and p and q even, then $K(N,p,q)$ is periodic of order two : it is invariant by a rotation of angle π around an axis S^1 and the linking number of its axis with the knot is equal to N*

Similarly to the Lissajous knots ([1]), these symmetries yields properties on the Arf invariant and Alexander polynomial that are described in the section.

We first have

Theorem 1 *Let N, p, q, ϕ be as above. Then the knot $K_{N,p,q,\phi}$ is strongly reversible.*

Proof Let us remind the reader that if θ is a real number and m is an integer, we have

$$\sin m(\theta+\pi) = (-1)^m \sin(m\theta), \ \cos m(\theta+\pi) = (-1)^m \cos(m\theta).$$

The change of parametrization $t \mapsto t+\frac{1}{2}$ doesn't change the knot as a whole ; it is induced by an ambiant symmetry of $\mathbb{S}^3$. We derive the following

Lemma 8 *The knots $K(N,p,q,0)$ and $K(N,p,q,\pi)$ are invariant under the diffeomorphism $\Phi_{N,p,q,}$ of $\mathbb{S}^3$ defined by*

$$(x,y,z,w) \mapsto ((-1)^N x, (-1)^N y, (-1)^q z, (-1)^p w).$$

It follows that the knots $K(N,p,q,0)$ and $K(N,p,q,\pi)$ - which are mirror images of one another - are reversible. This fact, together with Proposition 2 above, finishes the proof of the theorem. $\square$

According to the parities of the integers involved we can derive symmetries of the knot. First notice that $\Phi_{N,q,p}$ is an involution in all cases.

1. If N is even, then
    (a) if $p+q$ is even, $\Phi_{N,q,p}$ is an orientation preserving symmetry. From the non degeneracy condition (K is not singular), p and q can not be both even, hence p and q are odd ; then $\Phi_{N,q,p}$ is a rotation of π around an horizontal $\mathbb{S}^1$. Hence K is periodic of order two but in a weak sense since the linking number of the invariant axis of the rotation with K may be zero.

   (b) if $p+q$ is odd $\Phi_{N,q,p}$ is orientation reversing. and K is strongly fully amphichireal

2. If N is odd, then
   (a) if $p+q$ is even, $\Phi_{N,q,p}$ is orientation preserving. If p and q are both even, then $\Phi_{N,q,p}$ is a rotation of π around an vertical $\mathbb{S}^1$. its linking number with K is N. Hence K is periodic of order two. If p and q are both odd, then $\Phi_{N,q,p}$ is orientation preserving and has no fixed points.
   (b) if $p+q$ is odd $\Phi_{N,q,p}$ is orientation reversing. and K is strongly fully amphichireal

4.1 symmetries of Alexander polynomial

We can deduce from last considerations and [10] that

Corollary 2 *The simple minimal knot $K(N,p,q)$'s Alexander polynomial is a square modulo two, if p and q are not both odd, and the Arf invariant is then zero.*

4.2 symmetries of the braid

Using the braid description of the knots, these symmetries translate into symmetries of the braid. Let us write a point in the braid as (t,k) we mean the point $h_k(t)$ i.e. on the k-th strand. We distinguish two cases :

1. N **is even**. We notice that

$$\frac{2\pi}{N}(x+k)+\pi = \frac{2\pi}{N}(x+k+\frac{N}{2}) \tag{7}$$

   Thus, if p and q have the same (resp. a different) parity, $B_{N,q,p,0}$ is preserved (resp. transformed into its mirror image) by the following transformation :
   if $k \leq \frac{N}{2}$, $\quad (t,k) \mapsto (t, k+\frac{N}{2})$ if $k \geq \frac{N}{2}$, $\quad (t,k) \mapsto (t,k-\frac{N}{2})$.
   This symmetry switches the strands, while keeping the first coordinate fixed.



FIG. 2 – A knot that can not be simple minimal : 8_{17}

2. N **is odd**. We notice

$$\frac{2\pi}{N}(x+k)+\pi=\frac{2\pi}{N}(x+\frac{1}{2}+k+\frac{N-1}{2})=\frac{2\pi}{N}(x-\frac{1}{2}+k+\frac{N+1}{2}).$$

   If p and q have the same (resp. a different) parity, $B_{N,q,p,\phi}$ is preserved (resp. transformed into its mirror image) by the transformation
   if $k \leq \frac{N-1}{2}$ and $t \leq \frac{1}{2}$ then $(t,k) \mapsto (t+\frac{1}{2}, k+\frac{N-1}{2})$
   if $k \leq \frac{N-1}{2}$ and $t \geq \frac{1}{2}$ then $(t,k) \mapsto (t-\frac{1}{2}, k+\frac{N-1}{2})$
   if $k \geq \frac{N-1}{2}$ and $t \leq \frac{1}{2}$ then $(t,k) \mapsto (t+\frac{1}{2}, k-\frac{N+1}{2})$
   if $k \geq \frac{N-1}{2}$ and $t \geq \frac{1}{2}$ then $(t,k) \mapsto (t-\frac{1}{2}, k-\frac{N+1}{2})$
   This symmetry switches the first half of a strand with the second half of another.

5 A counterexample

Theorem 5 *A negative amphicheiral or chiral knot can not be the knot of a simple minimal or iterated minimal knot.*

Notice that the first candidate for a counterexample in the Rolfsen classification is the knot 8_{17} which is the first negative amphicheiral knot. This knot provides the first example of a prime knot which can not be a simple knot or iterated simple knot.

Proof. If the knot is simple, then theorem 1 shows that it can not be either chiral or negative amphicheiral.
Let us examine the case of an iterated minimal knot (of a simple knot) Suppose thus that there is an iterated minimal knot K that bounds a negative amphicheiral (or chiral) knot. Then K is contained in a tubular neighborhood of the companion knot, which is a simple minimal knot. The linking number κ of the satellite knot with a meridian of the tube is larger than two since the braid diagram of the satellite knot is identical to the braid diagram of an iterated torus knot. But the Alexander polynomial of K is a multiple of $P(x^{\kappa})$ where P is the Alexander polynomial of K which is impossible. $\square$

Notice though that it may be possible that these chiral knots are cable knots of minimal singular knots. This case requires a further investigation as we indicated in the introduction.

6 The algebraic crossing number

6.1 Definition - Background

We remind the reader that the *algebraic crossing number* of $B_{N,q,p,\phi}$ is the sum of the signs (i.e. $+1$ or -1) of its crossing points. We will denote it $e(B_{N,q,p,\phi})$. It is an invariant of the conjugation class of the braid but it is not an isotopy invariant of the knot $K_{N,q,p,\phi}$. However braids with three strands have been thoroughly studied by Birman and Menasco and we can derive from their work,

Theorem 2 *([B-M]). Let K (resp. K') be a knot represented by a 3-braid B (resp. B'). If K and K' are isotopic and K is neither trivial nor a $(2,k)$-link, then $e(B) = e(B')$.*

In the present case, where the braid comes from a branched immersion in 4-space, its algebraic crossing number can be seen as the number of double points which concentrate at the branch point. Namely

Theorem 3 *([Vi]) Let Σ be a closed Riemann surface without boundary, let M be an orientable 4-manifold and let $f : \Sigma \longrightarrow M$ be an embedding which has one branch point p. Suppose that in a neighbourhood of p, f is parametrized as in proposition 1, that it is a topological embedding in the* *neighbourhood of* p *and that the associated knot is* $K(N,p,q,\phi)$. *Then the degree of the normal bundle* Nf *is*

$$[f(\Sigma)].[f(\Sigma)] - e(B_{N,q,p,\phi})$$

where $[f(\Sigma)].[f(\Sigma)]$ *denotes the self-intersection number of* $f(\Sigma)$.

We immediately derive from Proposition 2 above that

Proposition 3 *Up to sign, the algebraic crossing number* $e(B(N,p,q,\phi))$ *does not depend on the phase* ϕ.

and that

Proposition 4 *Suppose* q *and* p *are of different parities. Then*

$$e(B(N,q,p,\phi)) = 0.$$

Remark A knot defined by a 2-stranded braid is either trivial or is a $(2,n)$ torus knot in which case the algebraic crossing number is n.

6.2 An estimate

The non-dependence on the phase allows us to prove

Proposition 5 *Let* N,q,p *as above and suppose that* p *and* q *are mutually prime. Then*

$$|e(B(N,q,p))| \leq N^2 + N - 4.$$

Remark This estimate does not depend on q : notice the sharp contrast with the case of the (N,q)-torus knot (i.e. when $p = q$) and every crossing number is positive : thus we have $e(B(N,q,q) = q(N-1)$.

Proof If k and l are different integers, $1 \leq k,l \leq N-1$, we denote by $M(k,l)$ the set of integers m such that the $t(m,k,l)$ given by the formula (*) above is a crossing point of the k-th and l-th strands.

Lemma 9 *The following correspondance is a bijection between* $M(k,l)$ *and* $M(N-1-k,N-1-l)$

$$t(m,k,l) \mapsto t(2q-m-1, N-1-k, N-1-l).$$

Proof. We notice that, if m verifies 1 w.r.t. the integers k and l, then $2q-m-1$ verifies

$$\frac{q}{N}(2N-(k+l)-2)-\frac{1}{2} \leq 2q-m-1 \leq \frac{q}{N}(2N-(k+l)-2)-\frac{1}{2}+\frac{2q}{N}.$$

In other words,

$$\frac{q}{N}((N-k-1)+(N-l-1))-\frac{1}{2} \leq 2q-m-1 \leq \frac{q}{N}(N-1-k)+(N-1-l))-\frac{1}{2}+\frac{2q}{N}.$$

□

Next, we go back to the expression of $R(.,\phi)$ above and see that we can choose a phase ϕ such that for every m, we have

$$R(m,\phi)=P([\frac{p}{q}m]).$$

At this point we introduce

$$\hat{M}=\{m \in M(k,l) \mid \frac{p}{q}m \in \mathbb{N}\}.$$

Since p and q are mutually prime, an element m of $\hat{M}$ is of the form $m=aq$, for some integer a.

Lemma 10 *If $m \in \hat{M} \cap M(k,l)$, then*
i) $m=q$
ii) $k+l=N-2$ or $k+l=N-1$ or $k+l=N$.
Conversely for every k,l verifying ii), $q \in M(k,l)$.

Proof The equality $m=aq$ yields

$$\frac{k+l}{N}-\frac{1}{2q} \leq a \leq \frac{k+l}{N}+\frac{2}{N}-\frac{1}{2q}.$$

Since $1 \leq k+l \leq 2N-3$, we derive that $a=1$. Thus $k+l$ satisfies

$$\frac{k+l}{N}-\frac{1}{2q} \leq 1 \leq \frac{k+l}{N}+\frac{2}{N}-\frac{1}{2q}.$$

Since $N>q$ the left hand-side yields $k+l \leq N$ and the right-hand side yields $k+l \geq N-2$.
We leave it to the reader to check the converse. □

Lemma 11 *If $m \neq q$, we have*

$$R(m,k,l,\phi) = R(2q-m, N-k-1, N-l-1, \phi) + 1.$$

Proof Since $\frac{p}{q}m$ is not an integer,

$$R(2q-m,\phi) = P([2p - \frac{p}{q}m]) = 1 + P([\frac{q}{p}m]) = 1 + R(m,\phi).$$

We denote by $\sigma(k,l)$ the signed number of intersection points of the k-th and l-th strands.
We have

$$\sigma(k,l)+\sigma(N-k-1,N-l-1) = -(-1)^{T(k,l)}\Sigma_{m\in M(k,l)}(-1)^{P(m)}\left((-1)^{R(m,\phi)} - (-1)^{R(2q-m-1,\phi)}\right).$$

If $k+l$ is not equal to either $N-2$, $N-1$ or N, then all m's in the sum above verify Lemma 10. Thus

$$|\sigma_{k,l} + \sigma(N-k-1, N-l-1)| \leq 2.$$

If $k+l = N$ (resp. $k+l = N-2$), then $(N-k-1)+(N-l-1) = N-2$ (resp. $(N-k-1)+(N-l-1) = N$).
Both pairs of strands $\{k,l\}$ and $\{N-k-1, N-l-1\}$ contain a crossing point for which $m = q$ and thus which does not verify Lemma 10. Thus

$$|\sigma_{k,l} + \sigma(N-k-1, N-l-1)| \leq 4.$$

Finally if $k+l = N-1$, then $\{k,l\} = \{N-k-1, N-l-1\}$. We have

$$2|\sigma(k,l) = |\sigma_{k,l} + \sigma(N-k-1, N-l-1)| \leq 4.$$

In order to put together these estimates, we point out the following :
if A is an integer, the number of pairs of strands $\{k,l\}$ such that $k+l = A$ is A. We derive the estimate

$$|e(b(N,p,q,\phi)| \leq 2\Sigma_{s=1}^{N-3}s + 4(N-2) + 2(N-1) = N^2 + N - 4.$$

7 Examples of Minimal Knots

Most of the examples of minimal knots given here can be described with the help of the Rolfsen table; i.e. they are knots that are the connected sum of prime knots with at most 10 minimum number of crosssings. In some cases however, we used the Hoste-Thistlewaite table provided in KnotTheory. It gives examples with a minimum crossing number reaching 16. We compute the Alexander Polynomial and Jones polynomial of the subdescribed minimal knots. In most cases we used the program KnotTheory to compute these polynomials as well as to draw the natural braids. Knots (in the Rolfsen table) were drawn using KnotPlot.

We will use Rolfsen's notation for the alexander polynomial :

$$[a_0 + a_1 + a_2 + \cdots + a_n] := x^n . \sum_{i=0}^{n} a_i \left(x + \frac{1}{x} \right)^i$$

7.1 General Properties

Beforehand we describe some useful properties of these knots.

As the simple knots are phase independant, the will be labelled as $K(N, p, q)$.

Lemma 4 *If*

$$\left\{ \begin{array}{l} x \equiv p \bmod N \\ x \equiv p \bmod 2q \end{array} \right. \tag{8}$$

then $K(N, x, q)$ *is isotopic to* $K(N, p, q)$

In particular, there is a $2qN$ periodicity for the appearance of knots with respect to p.

Lemma 5 *There is an infinite number of minimal representations of the trivial knot : for any* N *and* q, $K(N, N + q, q)$ *is isotopic to the trivial knot.*

Proof. Choose any two strands; we then can choose the phase such that the n labelling the crossings satisfies $0 \leq n < [\frac{2q}{N}]$. The crossing numbers appear then with the following sequence of signs as $+ + + + + + - - - - - - -$ where the numbers of plus and minus differ by one or zero. This means that the two strands can be deformed leaving the ends fixed such that their number of crossings is either one or zero without taking into account the other strands. But this is true for any two pairs of strands. Choose then strand 1. We can deform all others strands such that strand 1 is on top except for the strand that reaches the right end and that crosses strand 1 only once. We proceed similarly for all other strands. □

Lemma 6 *For all N, p, q, $K(N,p,q) = K(N,q,p)$*

Proof. We obtain $K(N,q,p)$ from $K(N,p,q)$ by adding the phase $\pi/2$ and by interchanging coordinates The invariance of the knot by a phase change yields the result. □.

We can check that

Lemma 7 *for any $N, q, a \in \mathbb{N}$, $K(N,aq,q)$ are isotopic to two type of knots : the torus knot $T(N,q)$ and a non trivial knot. Both have a braid group given by*

$$\left(\sigma_1^{\alpha_1} \circ \cdots \circ \sigma_{N-1}^{\alpha_{N-1}}\right)^N$$

In the case $T(N,q)$, all the $\alpha_i = +1$.

7.2 Torus Knots

It is known that Torus knots are not Lissajous knots [1]; this is in complete opposition with simple minimal knots :

Theorem 4 *All torus knot are minimal knots : $T(a,b)$ can be realized as $K(a,b,b)$. Moreover the regular knots $K(2,p,q)$ is trivial if $p \neq q$; else it is the torus knot $T(q,2)$.*

The first part is direct; the second part is a consequence of last section on the algebraic number of minimal braids with two strands.

7.3 Minimal Knots with an odd number of strands

We will consider first knots whose minimal braid has three strands. The braid representative of $K(3,p,q)$ is $\prod_{i=1}^{q} \sigma_1^{\alpha_i}\sigma_2^{\beta_i}$, $\alpha_i, \beta_i = \pm 1$ and σ_i denotes the ith crossing ordered from left to right on the braid diagram; strands are numbered top down. If the first two crossings of the braid corresponds respectively to the crossings of the couple of strands $(1,2)$ and $(1,3)$ then the k-th pairs of crossings correspond to the intersection of the couple of strands $(\tau^k(1), \tau^k(2))$ and $(\tau^k(1), \tau^k(3))$ where τ is the cyclic permutation $(1,2,3)$.

7.3.1 K(3, . , 4)

The number of crossings is at most $q.(N-1) = 8$; hence, if theses knots are prime, they must be in the Rolfsen table; If they are not prime then they are connected sums of knots in the Rolfsen table. We describe in this paragraph all such knots. Only five knots appear periodically as p varies; From lemma 4 there is a global 24-periodicity with respect to p but there are also other symmetries that are not accounted for.

1. The torus knot $T(3,4) = K(3,4,4p) = K(3,20,4) \cdots$ This case generalizes to all $K(N,p,p)$ or $K(N,ap,p)$ for suitable a. These knots are clearly reversible.
2. The square knot $3_1\#\bar{3}_1 = K(3,5,4) = K(3,4,5)$. This is the first example in the literature of a minimal knot that is not toric (cf. [MW]). In a way, this the smallest knot with respect to the lexicographic ordering on N, p, q. It is fully amphicheiral and $P = [-1+1]^2$.
3. The trivial knot $K(3,7,4) = 1$. This is knot of type $K(N, N+q, q)$ which are all trivial. Notice that $K(3,k,4) = 1$ for $k = 5, 11, 13, 19...$
4. The connected sum of the eight knot and the square knot is $4_1\#(3_1\#\bar{3}_1) = K(3,8,4)$. ($p = 8, 16...$). This is a knot of type $K(N, aq, q)$; it is either the torus knot $T(N,p)$ or a knot whose braid group is $\left(\sigma_1\sigma_2^{-1}\right)^4$. This knot is fully amphicheiral
5. The first non toric and non trivial prime knot, the eight knot $4_1 = K(3,10,4) = K(3,10,4) = K(3,22,4)...$; it is fully amphicheiral. $P = [-3+1]$.

7.3.2 K(3, . , 5)

The number of crossings is at most 10 which still allows us to check in the Rolfsen table. Only four knots appear periodically as p varies. We just write down the smallest p for each knot.

1. The torus knot $T(3,5)$. realized by $K(3,5,5)$.
2. The prime knot $10_{155} = K(3,7,5)$; it is reversible. As in the case of Lissajous knot the appearance of this knot for the values $3, 7, 5$ suggests that it is difficult to find a topological caracterization of these minimal knots. This is even clearer in the case $K(4,11,5)$ described below.
3. The trivial knot $K(3,8,4) = 1$.



Fig. 3 – minimal braid and knot of K(3,5,4)



Fig. 4 – One of many minimal representation of a trivial knot : K(3,7,4)



FIG. 5 – The eight knot realized as K(3,10,4)

4. The connected sum $6_2\#\bar{6}_2 = K(3,10,5)$. This is a knot of type $K(N,aq,q)$; $(= K(3,20,5)$, but $K(3,25,5) = K(3,35,5)$ is the torus knot $T(3,5)$ and $K(3,10,5) = K(3,40,5)$...)

7.3.3 K(3, . , 7)

The number of minimal crossings is at most 14, two of them have a minimal crossing number of 14 ; to check all cases we need consult the Hoste-Thistlewaite table. Six different knots appear periodically.

1. The torus knot $K(3,3,7) = T(3,7)$.
2. The knot $K(3,8,7)$ is fully amphicheiral and $P = [1-1+1]^2$.
3. The trivial knot $K(3,10,5) = 1$.
4. The prime knot $14N27120 = K(3,11,7)$ which is reversible ! This is the first occurence of a non amphicheiral knot with odd numbers , N, p, q which is not a torus. $P = [7-5+3-1]$.
5. The connected sum $K(3,14,7)$; it is amphicheiral, and $P = [7-6+4-1]^2$. This is a knot of type $K(N,aq,q)$, hence its braid group is $\left(\sigma_1^{-1}\sigma_2\right)^7$
6. The prime knot $K(3,19,7) = 14N11995$; it is reversible and $P = [7-5+3-1]$.



Fig. 6 – An example of a torus knot : K(3,5,5)



Fig. 7 – minimal braid of $10_{155} = K(3,7,5)$



FIG. 8 – minimal braid of $K(3,10,5) = (6_2 \# \bar{6}_2)$ and knot 6_2



FIG. 9 – minimal braids of $K(3,11,7) = 14N27120$ and $K(3,14,7)$



FIG. 10 – minimal braids of $K(3,19,7) = 14N11995$ and $K(5,16,6) = 14N17954$

With $N = 3, q \geq 8$ the identification becomes very difficult since the number of crossings is a priori bigger than 16! We conclude this paragraph with an example of a minimal knot with $N = 5$ i.e. whose minimal braid representation has 5 strands.

7.3.4 K(5,22,6)

We will consider knots whose minimal braid has five strands. The braid representative of $K(5,p,q)$ is $\prod_{i=1}^{q} \sigma_2^{\alpha_i}\sigma_4^{\beta_i}\sigma_1^{\gamma_i}\sigma_3^{\delta_i}$, $\alpha_i, \beta_i, \gamma_i, \delta_i = \pm 1$. The first four crossings of the braid on the strands corresponds to the couples $(2,3), (4,5), (1,3), (2,5)$ and the k-th quadruple corresponds to the crossings $\tau^k(a,b)$ where $\tau = (1,3,5,4,2)$.

We will only describe an example of prime minimal knot. In each case we need to check on the braid representations that the minimal crossing number is less than 16. We try to minimize the number of crossings on the minimal braid for each knot such that the crossing number is less than 15. We will number the strands top down according to the order of the strands on the far left of the braid.

1. $K(5,22,6) = 7_7$ (fig. 11). Consider the minimal braid of this knot; flip to the bottom the first "hill" of the strand 4, we reduce the crossing number by 4; Flip the first valley of strand 2 to the top , we can reduce the crossing number by 4. Then moving downwards the first hill of



FIG. 11 – minimal braid of knot $K(5, 22, 6) = 7_7$

strand 3 reduces the number by 2; hence it suffices to check in the table of knots with less than 14 crossings.

It seems that the Alexander polynomial of all the examples computed have a highest order coefficient equal to plus or minus one; It may be that all these simple minimal knots are fibered when N is odd. hence we conjecture :

Conjecture 1 *Simple minimal knots with an odd number of strands are fibered.*

This is not the case when N is even as we shall see now.

7.4 Minimal Knots with an even number of strands

We will only discuss the case $N = 4$, the case $N = 2$ has been dealt with in the computation of its algebraic crossing number in section 4.

7.4.1 K(4, . 5)

We will consider first knots whose minimal braid has four strands. The braid group of $K(4, p, q)$ is $\prod_{i=1}^{q} \sigma_1^{\alpha_i} \sigma_3^{\beta_i} \sigma_2^{\gamma_i}$, $\alpha_i, \beta_i, \gamma_i = \pm 1$. If the first three crossings of the braid corresponds respectively to the crossings of the couple of strands $(1, 2), (3, 4), (1, 4)$, the k-th triple corresponds to the image of the first three by the permutation τ^k where $\tau = (1, 2, 4, 3)$



Fig. 12 – The first non-fibered prime knot : K(4,13,5)

The number of crossings is at most 15 Six different knots appear periodically.

1. The torus knot $T(4,5)$.
2. The sum of two prime knots $K(4,7,5)$. it is reversible and $P = [17 - 12 + 4]$. As the highest coefficient of the Alexander polynomial is 4, this knot is not fibered ; it provides the first occurence of a non-fibered minimal knot.
3. The trivial knot $K(4,9,5) = 1$.
4. The prime knot $15N166131 = K(4,11,5)$ is reversible and $P = [37 - 28 + 12 - 2]$.
5. The non fibered prime knot $9_{46} = K(4,13,5)$, reversible and $P = [5 - 2]$. In fact we can reduce easily the crossing number of the minimal braid to 13 which makes computations easier.

Corollary 1 *Singularity knots of minimal branch points are not necessarily fibered*

Marc Soret at marc.soret@lmpt.univ-tours.fr
Université F. Rabelais, Dép. de Mathématiques, 37000 Tours, France
Marina Ville at mville@math.jussieu.fr
C.N.R.S. & Dept of Mathematics,
Northeastern University, 360 Huntington Avenue, Boston MA 02115, USA

Références

[1] M. Bogle, J. Hearst, V. Jones & L. Stoilov LISSAJOUS KNOTS , Journal of Knot Theory and Its Ramifications, Vol.3, No. 2 (June 1994), pp. 121-140

[2] E. Brieskorn & H. Knorrer PLANE ALGEBRAIC CURVES , Birkhauser, 1986.

[3] J. Birman & W. Menasco STUDYING LINKS VIA CLOSED BRAIDS III : CLASSIFYING LINKS THAT ARE CLOSED 3-BRAIDS, Pac. Jour. Maths 161, 25-115 (1993)

[4] G. Burde & H. Zieschang KNOTS Walter de Gruyter, 1985

[5] R. Hartley & A. Kawauchi POLYNOMIALS OF AMPHICHEIRAL KNOTS, Math. Ann. 243, 63-70 (1979)

[6] V. Jones & J. Przytycki LISSAJOUS KNOTS AND BILLIARD KNOTS, Banach Center Publications 42, 145-163 (1998).

[7] Mathematica Package KNOTTHEORY,
$http://katlas.math.toronto.edu/wiki/Main_Page$

[8] R. Sharein KNOTPLOT, $http://www.pims.math.ca/knotplot$

[9] J. Milnor SINGULAR POINTS OF COMPLEX HYPERSURFACES Ann. of Math. Studies 61, (1968) Princeton University Press,

[10] K. Murasugi ON PERIODIC KNOTS , Comment. Math. Helv. 46 (1971), 162-174.

[11] Micallef & B. White THE STRUCTURE OF BRANCH POINTS IN MINIMAL SURFACES AND IN PSEUDOHOLOMORPHIC CURVES ANNALS OF MATHS, 139, (1994), 35-85

[12] D. Rolfsen KNOTS AND LINKS Mathematical Lecture Series, 7, Publish or Perish, 1990

[13] M. Ville BRANCHED IMMERSIONS AND BRAIDS, preprint (2006).